



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

*Sanitary report from Santiago.*SANTIAGO DE CUBA, *August 1, 1896.*

SIR: I have the honor to inform you that 108 deaths have taken place at Santiago de Cuba during the week ended Saturday, August 1. Of these, 21 were from yellow fever, 39 from smallpox, 1 from typhomalaria, 5 from remittent fever, 6 from enterocolitis, and the rest from common diseases, both acute and chronic, and of noncontagious character.

Yellow fever has augmented greatly this week, and there are quite a number of cases at the military hospital at present. Reports from different parts of the island state that yellow fever is prevalent among the soldiers, and as the hot season is now at its culminating point I am afraid an epidemic of this terrible malady will be established.

Smallpox is decreasing somewhat and the cases presenting a milder character. Malaria is very deadly, the remittent form of the fever prevailing. Diarrhea under the form of enterocolitis, is still causing many victims among the infantile population. Temperature very high, alternating with heavy showers of rain.

Respectfully,

HENRY S. CAMINERO,
Sanitary Inspector, U. S. M. H. S.

*Report on yellow fever and other infectious diseases.*HABANA, CUBA, *August 1, 1896.*

SIR: I have the honor to inform you that there were 1,062 deaths in this city during the month of July, 1896. One hundred and twenty-two of these deaths were caused by yellow fever, of which 97 occurred in the military hospital, 60 were caused by enteric fever, 22 by so-called pernicious fever, 11 by paludal fever, 2 by diphtheria, 27 by dysentery, 83 by enteritis, 84 by smallpox, 10 by measles, 14 by pneumonia, 3 by glanders, and 148 by tuberculosis.

The newspaper "Diario de la Marina," published to-day the following: "In Yaguajay, Caibarien Remedios, Camajuani, Vuelta, Zulueta, Buena Vista, and Placetas, towns of Santa Clara Province, district of Remedios, yellow fever has taken an epidemic form and attacks natives coming from the interior of the island to seek protection in these crowded places."

Report of the Spanish army sanitary corps for the first ten days of July, 1896, including all hospitals of the Island of Cuba.

	Existing July 1.	New cases.	Recovered.	Deaths.	Remaining July 10.
Yellow fever.....	792	911	474	247	982
Dysentery	146	53	53	3	143
Malaria	927	208	246	2	885
Enteric fever.....	72	28	21	9	70
Wounded	728	133	144	5	712
General diseases.....	8, 076	5, 769	5, 066	327	8, 452

The smallpox is not mentioned in the report.

Very respectfully,

MANUEL S. CASTELLANOS,
Sanitary Inspector, U. S. M. H. S.